

**CORRIGENDIUM: KRULL DIMENSION, OVERRINGS AND
SEMISTAR OPERATIONS OF AN INTEGRAL DOMAIN. [J.
ALGEBRA, 321 (2009), 1497–1509]**


A. MIMOUNI

Theorem 2.9 Let R be an integrally closed domain. Then the following conditions are equivalent.

- (i) $|O(R)| = 7 + \dim R$.
- (ii) $|SSFc(R)| = 7 + \dim R$.
- (iii) R is a Prüfer domain, and either
 - (a) R has exactly three maximal ideals M_1, M_2, M_3 and $\text{Spec}(R) = \{0 = P_0 \subsetneq P_1 \subsetneq \cdots \subsetneq P_{r-1} \subsetneq M_1 \cap M_2 \cap M_3\}$; or
 - (b) $\dim R \geq 5$, R has exactly two maximal ideals M and N and $\text{Spec}(R) = \{0 = P_0 \subsetneq P_1 \subsetneq \cdots \subsetneq P_{r-1} \subsetneq P_r = M, N, P_{r-4} \not\subseteq N, P_{r-5} \subsetneq N\}$; or
 - (c) $\dim R \geq 2$, R has exactly two maximal ideals M and N and $\text{Spec}(R) = \{0 = P_0 \subsetneq P_1 \subsetneq \cdots \subsetneq P_{r-2} \subsetneq P_{r-1} \subsetneq P_r = M, P_{r-2} \subsetneq Q \subsetneq N\}$.

Proof. Only Case 1 need a slight modification proof.

Case 1 $|Max(R)| = 3$. Assume that $Max(R) = \{M_1, M_2, M_3\}$ and set $ht M_1 = \dim R = r \geq 2$. Let $0 = P_0 \subsetneq P \subsetneq M_1$ be a chain of prime ideals of R . We claim that $P_{r-1} \subseteq M_2 \cap M_3$. Indeed, suppose that $P_{r-1} \not\subseteq M_2$. Then $O(R) \supseteq \{R, L, R_{P_1}, \dots, R_{P_{r-1}}, R_{M_1}, R_{M_2}, R_{M_3}, R_{M_1} \cap R_{M_2}, R_{M_1} \cap R_{M_3}, R_{M_2} \cap R_{M_3}, R_{P_{r-1}} \cap R_{M_2}\}$. Hence $7 + r = 7 + \dim R = |O(R)| \geq 9 + r = 9 + \dim R$, which is absurd. Thus the assertions (a) is satisfied and $\text{spec}(R)$ is of the form.



□

Example 3.4. A one-dimensional Noetherian local domain R such $|O(R)| = 3 + \dim R$ and $|SSFc(R)| = 5 + \dim R$.

2000 *Mathematics Subject Classification.* Primary 13G05, 13A15, 13F05; Secondary 13G05, 13F30.

Key words and phrases. semistar operation, Krull dimension, Prüfer domain, *fgv* domain.
This work is supported by KFUPM.

Let k be a field and X an indeterminate over k . Set $R = k[[X^3, X^4, X^5]]$. Then R is a one-dimensional Noetherian local domain which is not divisorial (Lemma 3.1, since $M^{-1} = k[[X]]$ and $R \subsetneq k[[X^2, X^3]] \subsetneq M^{-1}$). Also it is easy to see that $O(R) = \{R \subsetneq R_1 = k[[X^2, X^3]] \subsetneq R_2 = R' = k[[X]] \subsetneq L = qf(R) = k((X))\}$ and each proper overring of R is divisorial (Lemma 3.1) so fgv domain. Clearly $|O(R)| = 4 = 3 + \dim R$, however $|SSFc(R)| = 6 = 5 + \dim R$ (in fact $SSFc(R) = \{e, \bar{d}, \overline{*_{R_1}}, \bar{t}, \{*_R\}_{1 \leq i \leq 2}\}$, where $E^{\overline{*_{R_1}}} = ER_1 \cap E_v$).

Example 3.5. A one-dimensional Noetherian local domain with $|O(R)| = |SSFc(R)| = 5 + \dim R$.

Let k be a field and X an indeterminate over k . Set $R = k[[X^2, X^9]]$. Then R is a one-dimensional Noetherian local domain which is divisorial (since $M^{-1} = k[[X^2, X^7]]$). Also it is easy to see that $O(R) = \{R \subsetneq R_1 = k[[X^2, X^7]] \subsetneq R_2 = k[[X^2, X^5]] \subsetneq R_3 = k[[X^2, X^3]] \subsetneq R_4 = R' = k[[X]] \subsetneq L = qf(R) = k((X))\}$ and each overring of R is divisorial (Lemma 3.1) so fgv domain. By Theorem 2.3, $|SSFc(R)| = |O(R)| = 6 = 5 + \dim R$ (in fact $SSFc(R) = \{e, \bar{d}, \{*_R\}_{1 \leq i \leq 4}\}$).

Example 3.6. The following is an example of a domain R with $|O(R)| = 5 + \dim R$ and $|SSFc(R)| = \infty$.

Let $\mathbb{Q}$ be the field of rational numbers, X an indeterminate over $\mathbb{Q}$. Set $V = \mathbb{Q}(\sqrt{2}, \sqrt{3})[[X]] = \mathbb{Q}(\sqrt{2}, \sqrt{3}) + M$, where $M = XV$ and $R = \mathbb{Q} + M$. By [2, Theorem 2.1], R is a one-dimensional Noetherian local domain with maximal ideal M . Since each overring of R is comparable to V ([2, Theorem 3.1]), then it is easy to see that $O(R) = \{R, V, L = qf(R), R_1 = \mathbb{Q}(\sqrt{2}) + M, R_2 = \mathbb{Q}(\sqrt{3}) + M, R_3 = \mathbb{Q}(\sqrt{6}) + M\}$ and all proper overrings of R are divisorial ([2, Corollary 4.4]). Hence $|O(R)| = 6 = 5 + \dim R$. Now, since $[\mathbb{Q}(\sqrt{2}, \sqrt{3}) : \mathbb{Q}] = 4$ and Q is infinite, $|S(R)| = \infty$ and therefore $|SSFc(R)| = \infty$.

Example 3.7. A one-dimensional Noetherian local domain R such that $|SSFc(R)| = |O(R)| = 6 + \dim R$.

Let k be a field and X an indeterminate over k . Let $R = k[[X^2, X^{11}]]$. Since $M^{-1} = k[[X^2, X^9]]$, by Lemma 3.1, R is divisorial. Now, it is easy to see that $O(R) = \{R \subsetneq R_1 = k[[X^2, X^9]] \subseteq R_2 = k[[X^2, X^7]] \subsetneq R_3 = k[[X^2, X^5]] \subsetneq R_4 = k[[X^2, X^3]] \subsetneq R_5 = R' = k[[X]] \subsetneq L = qf(R) = k((X))\}$. Also it is easy to check that each overring is divisorial (Lemma 3.1). By Theorem 2.3, $|SSFc(R)| = |O(R)| = 7 = 6 + \dim R$.

Example 3.9. Just change $R_1 = k[[X^3, X^5]]$ to $R_1 = k[[X^2, X^9]]$.

DEPARTMENT OF MATHEMATICS AND STATISTICS, KING FAHD UNIVERSITY OF PETROLEUM & MINERALS, P. O. BOX 278, DHAHRAN 31261, SAUDI ARABIA
E-mail address: amimouni@kfupm.edu.sa